

LEGIONNAIRES' DISEASE



FORTBEND
HEALTH & HUMAN SERVICES
Prevent. Promote. Protect.

Legionnaires' disease, a type of severe pneumonia, is caused by breathing in small droplets of water that contain Legionella.



Signs and symptoms of Legionnaires' disease

Cough

Fever

Headache

Muscle Aches

Shortness of Breath



Legionnaires' disease is serious, but can be treated with antibiotics.

Legionnaires' disease is not contagious and can be treated with antibiotics (*drugs that kill bacteria in the body*) when caught early.

Source: www.cdc.gov/legionella

Disclaimer: This information does not constitute medical advice or a doctor-patient relationship

Certain people are at increased risk for Legionnaires' Disease.

Most healthy people do not get Legionnaires' disease after being exposed to Legionella. However, certain factors can increase your chance of getting sick. These risk factors include:



Being 50 years or older



Having a weakened immune system



Being a current or former smoker



Taking medication that weakens your immune system



Having chronic lung disease

Common sources of infection outbreaks of Legionnaires' disease

Sources of infection outbreaks of Legionnaires' disease are often associated with large water systems, like those found in hospitals, hotels, and cruise ships, specifically:



**Water used for showering
(potable water)**



Hot tubs



Decorative fountains



**Cooling towers
(parts of large air conditioning systems)**

Source: www.cdc.gov/legionella

Disclaimer: This information does not constitute medical advice or a doctor-patient relationship

LEGIONELOSIS

La enfermedad del legionario, un tipo de neumonía grave causada por inhalar pequeñas gotitas de agua que contiene Legionella.



Signos y síntomas de la enfermedad del legionario



Tos



Fiebre



Dolor de cabeza



Dolores musculares



Dificultad para respirar




La enfermedad del legionario es grave, pero se puede tratar con antibióticos.

La enfermedad del legionario no es contagiosa y se puede tratar con antibióticos (medicamentos que matan las bacterias en el cuerpo) cuando se detecta a tiempo.

Fuente: www.cdc.gov/legionella

Descargo de responsabilidad: Esta información no constituye consejo médico ni una relación médico-paciente



Ciertas personas tienen un riesgo más alto de contraer la enfermedad del legionario.

La mayoría de las personas sanas no contraen la enfermedad del legionario después de estar expuestas a Legionella. Sin embargo, ciertos factores de riesgo pueden aumentar su probabilidad de enfermarse. Estos factores de riesgo incluyen:



Tener 50 años o más



Tener un sistema inmunológico debilitado



Ser fumador o ex fumador



Tomar medicamentos que debilitan su sistema inmunológico



Tener una enfermedad pulmonar crónica

Fuentes comunes de brotes de infección de la enfermedad del legionario

Los brotes de infección de la enfermedad del legionario a menudo se asocian a sistemas de agua grandes, como los que se encuentran en hospitales, hoteles y cruceros, específicamente:



Agua utilizada para ducharse (agua potable)



Bañeras para hidromasaje o jacuzzi



Fuentes decorativas



Torres de refrigeración

Fuente: www.cdc.gov/legionella

Descargo de responsabilidad: Esta información no constituye consejo médico ni una relación médico-paciente